

B_s^0 mixing at D0 experiment

T. Moulik (for the D0 collaboration)

Department of Physics, University of Kansas, Lawrence, KS 66045, USA

In this report, we present a preliminary measurement of the B_s^0 mixing parameter using samples of four partially reconstructed semileptonic B_s^0 decays and one fully reconstructed hadronic decay mode, corresponding to approximately 2.4 fb^{-1} of integrated luminosity. We perform an unbinned likelihood fit to the proper decay length and obtain $\Delta m_s = 18.53 \pm 0.93(\text{stat.}) \pm 0.30(\text{syst.}) \text{ ps}^{-1}$.

1. INTRODUCTION

Particle-antiparticle oscillations are observed and well established in the B_d^0 system. The mass difference Δm_d is measured to be $\Delta m_d = 0.507 \pm 0.005 \text{ ps}^{-1}$ [1]. B_s^0 mesons are known to oscillate with a high frequency according to standard model predictions. Observing the oscillations in the B_s^0 system has been an important focus of the B physics program at both the D0 and the CDF experiments at Tevatron. D0 reported direct limits on the B_s^0 mixing parameter Δm_s [2] using the $B_s^0 \rightarrow D_s^- \mu^+ \nu X$, $D_s^- \rightarrow \phi \pi^-$ decay mode ¹. CDF reported a measurement of this parameter exceeding 5σ significance [3]. The measurement of this parameter is an important test of the CKM (Cabibbo Kobayashi Maskawa) formalism of the standard model, and combining it with a measurement of Δm_d allows us to reduce the error on V_{td} and constrain one side of the CKM triangle. This report describes the measurement of the B_s^0 mixing parameter at the D0 experiment using 2.4 fb^{-1} of data with 3σ significance.

We use the central tracker, muon chambers and calorimeters to reconstruct the B decays. Details of the detector can be found elsewhere [4]. We use a single inclusive muon trigger or a di-muon trigger to accumulate samples for B_s^0 mixing studies. In the case of di-muon trigger the other muon acts as the tag muon used to identify the flavor of the B meson which we discuss later. The trigger requires a good muon identified by the muon chamber with a matching track in the central tracker in the pseudo-rapidity range of $|\eta| < 2.0$. The triggers use p_T cuts between $3 - 5 \text{ GeV}/c$ and the trigger is prescaled or turned off depending on the luminosity. Hence for the $B_s^0 \rightarrow D_s^- e^+ \nu X$ decay mode, and the hadronic decay mode, we are using a tagged sample with the muon acting as a tag.

2. B_s^0 DECAYS SAMPLE SELECTION AND RECONSTRUCTION

D0 reported direct limits on B_s^0 oscillations using the decay mode $B_s^0 \rightarrow D_s^- \mu^+ \nu X$ decays with $D_s^- \rightarrow \phi \pi^-$ [2]. In this report, we present a preliminary measurement of the B_s^0 oscillations [5], using four additional decay modes. We use three additional semileptonic decay modes namely $B_s^0 \rightarrow D_s^- e^+ \nu X$, with $D_s^- \rightarrow \phi \pi^-$ and $B_s^0 \rightarrow D_s^- \mu^+ \nu X$ decays with $D_s^- \rightarrow K^{*0} K^-$, $K^{*0} \rightarrow K^+ \pi^-$ (The K^{*0} and the ϕ candidates are required to be consistent with known mass and width [1] of these two resonances) and $D_s^- \rightarrow K_S K^-$. We also use a fully reconstructed hadronic mode, namely, $B_s^0 \rightarrow D_s^- \pi^+ + X$ decay, with $D_s^- \rightarrow \phi \pi^-$.

B_s^0 decays are identified in their semileptonic modes, both in muon and electron mode. Muons are required to have $p_T > 1.5 \text{ GeV}/c$, and to be in the pseudo-rapidity region of $|\eta| < 2.0$. Electrons are required to have $p_T > 2.0 \text{ GeV}/c$ and are identified in a pseudo-rapidity region of $|\eta| < 1$. The D_s^- and B_s^0 decay products are constrained to originate from a common vertex and the B_s^0 and D_s^- decay vertices are required to be significantly displaced from the $p\bar{p}$ collision vertex.

¹Charge conjugated states are implied throughout the text

The $K^{*0} \rightarrow K^+\pi^-$ decay mode requires special treatment on account of large reflections, as both real physics processes and combinatorial background contribute to the signal peak. The peak is comprised of the signal mode, $D_s^- \rightarrow K^{*0}K^-$, $K^{*0} \rightarrow K^+\pi^-$, the physics processes, $D^+ \rightarrow K^-\pi^+\pi^+$ or $D^+ \rightarrow K^{*0}\pi^+(K^{*0} \rightarrow K^+\pi^-)$, $\Lambda_c^+ \rightarrow K^+\pi^-p^+$, $D^+ \rightarrow K^{*0}K^+(K^{*0} \rightarrow K^+\pi^-)$ (Cabibbo suppressed) and combinatorial background. We fit for these contributions in an unbinned likelihood fit. Fig. 1 shows the fit to the mass distribution of the $M_{K\pi K}$ system with the individual contributions superimposed.




Figure 1: The $M_{(K\pi)K}$ invariant mass distribution where $D_s \rightarrow K^{*0}K$ for $B_s^0 \rightarrow D_s^- \mu^+ \nu X$ decay mode

3. FLAVOR TAGGING

The flavor of the initial state of the B_s^0 is determined using a likelihood ratio method, based on the properties of the other b-hadron in the event (opposite side tagging) and the properties of the particles produced in association with the reconstructed B_s^0 meson (same side tagging) and we then combine the OST and the SST taggers. The performance of any tagger is characterized by its efficiency defined as $\varepsilon = N_{tag}/N_{tot}$ (where N_{tag} is the number of tagged B_s^0 mesons and N_{tot} is the total reconstructed B_s^0 mesons), and the dilution $\mathcal{D}$ which is defined as $\mathcal{D} = \frac{N_{RS} - N_{WS}}{N_{tag}}$ (where N_{RS} and N_{WS} are the right-sign and wrong-sign tags respectively).

The OST is calibrated using $B \rightarrow \mu + \nu D^{*-}$ data events, and we obtain the dilution $\mathcal{D}$ as a function of a tag variable d (whose sign indicates a b or $\bar{b}$, and its value indicates the “ b ”-ness of the tag), to provide an event-by-event “predicted” dilution which is used in the unbinned likelihood fit described in section 4. More details on the development of the OST and results can be found in [6]. The total OST effective tagging power of $\varepsilon \mathcal{D}^2 = (2.48 \pm 0.21_{-0.06}^{+0.08})\%$.

The combined tagger is calibrated using decay modes, $B_s^0 \rightarrow J/\psi \phi$ (using both data and MC) and $B_s^0 \rightarrow \mu D_s(\phi\pi)$ (using MC only) (See Fig. 2). The total combined tagging power is $\varepsilon \mathcal{D}^2 = 4.49 \pm 0.88\%$. More details on the SST and the combined tagger can be found in [7].

The SST is used by $B_s^0 \rightarrow D_s^- \mu^+ \nu X$, with $D_s^- \rightarrow \phi \pi^-$ decay mode only. All the other decay modes use the OST.

4. UNBINNED LIKELIHOOD FIT AND AMPLITUDE SCAN

4.1. Visible Proper decay length and scale factor

In semileptonic decays the proper time gets smeared due to the presence of neutrino. To take this into account we introduce a K factor estimated from Monte Carlo (MC) simulation. It is defined as $K = p_T(lD_s)/p_T(B)$. The real




Figure 2: The Combined Tagger dilution calibration

proper decay length is related to the measured or visible proper decay length (l_M), by the relation $ct(B_s^0) = l_M K$ where $l_M = M(B_s^0) \cdot (L_T)/(p_T(lD_s))$ is the measured visible proper decay length (l_M). L_T is the distance from the primary vertex to the B_s^0 decay vertex in the transverse plane projected onto the lD_s momentum, and $M(B_s^0)$ is the mass of the B_s^0 meson as obtained from PDG [1].

The l_M uncertainty is determined by the vertex fitting procedure, track parameters, and track parameter uncertainties. To account for any imperfections in modeling of detector uncertainties, we use scale factors that are dependent on the topology of hits in the silicon micro-strip tracker to correct the l_M uncertainty. These scale factor corrections were obtained from QCD data and MC samples.

4.2. Likelihood Fit

An unbinned likelihood fit is used to describe and fit for the B_s^0 oscillation. All flavor tagged events with $1.72 < M(D_s) < 2.22 \text{ GeV}/c^2$ are used in the fit. The probability for oscillated (osc) and non-oscillated (nos) B_s^0 decays, as a function of true proper decay length, x , can be written as

$$p_s^{nos}(x) = \frac{K}{c\tau_{B_s}} \cdot e^{-\frac{K}{c\tau_{B_s}}} \cdot 0.5 \cdot (1 + \mathcal{D} \cos(\Delta m_s \cdot Kx/c)) \quad (1)$$

$$p_s^{osc}(x) = \frac{K}{c\tau_{B_s}} \cdot e^{-\frac{K}{c\tau_{B_s}}} \cdot 0.5 \cdot (1 - \mathcal{D} \cos(\Delta m_s \cdot Kx/c)) \quad (2)$$

The sample is mostly composed of B_s^0 decays with some contributions coming from B_d and B_u mesons also. Similar equations would hold for these decay modes with the oscillatory term getting modified appropriately.

Taking into account momentum uncertainty, and convoluting over detector resolution, the probability for the j -th decay mode, as a function of the measured proper decay length, can be written as below :

$$p_j^{nos/osc}(x^M) = \sum_j f_j(K) dK \epsilon_j(x^M) \int dx g(x - x^M, x) p_j^{nos/osc}(x, K) \quad (3)$$

$f_j(K)$ is the K-factor distribution and we sum over the K -factor bins. $\epsilon_j(x^M)$ is the reconstruction efficiency as a function of lifetime cuts. The fraction of each decay mode contribution is calculated from Monte Carlo. A similar

probability equation holds for the combinatorial background (p_{bg}). The background contribution comes from prompt decays with the $D_s^- \mu(e)$ vertex coinciding with the primary vertex, background with fake vertices distributed around the primary vertex, and long lived background.

The likelihood can then be written as follows :

$$\mathcal{L} = \prod_{i=1}^N ((1 - f_{sig})p_{i,bg} + f_{sig}p_{i,sig}) \quad (4)$$

The amplitude method was first proposed elsewhere [8]. It involves modifying the likelihood term by introducing an amplitude term $\mathcal{A}$ in front of the oscillatory cosine term, such that,

$$\mathcal{L} \propto 1 \pm \mathcal{A} \mathcal{D} \cos(\Delta m_s t) \quad (5)$$

We minimize $-2\ln\mathcal{L}$ fitting for the parameter $\mathcal{A}$ while $\mathcal{D}$ is known and Δm_s is varied. The value of Δm_s where $\mathcal{A}$ is consistent with 1 and inconsistent with 0 then gives the measurement of the Δm_s parameter. All values of Δm_s for which $\mathcal{A} + 1.645 \sigma_{\mathcal{A}} < 1$ are excluded at 95% confidence level. The sensitivity of the mixing measurement is defined as the Δm_s value for which $1.645 \sigma_{\mathcal{A}} = 1$.

The combined amplitude scan and the difference between log-likelihood at minimum and elsewhere ($\Delta\ln\mathcal{L}$), as a function of Δm_s , can be seen in Fig. 3. $\Delta\ln\mathcal{L}$ is derived from the amplitude scan. More details can be found in [8].




Figure 3: The combined amplitude and likelihood scan.

From a parabolic fit around the minimum of the $\Delta\ln\mathcal{L}$ distribution, we obtain $\Delta m_s = 18.53$ with statistical error of 0.93 ps^{-1} . The systematic error is estimated as 0.3 ps^{-1} , thus giving us a preliminary measurement of $\Delta m_s = 18.53 \pm 0.98 \text{ ps}^{-1}$ (stat.+syst.). From the depth of the likelihood minima, we calculate an overall significance of this measurement as 2.9σ .

References

- [1] W. M. Yao *et al.*, Particle Data Group, J. Phys. G**33** (2006)
- [2] V.M. Abazov *et al.*, D0 Collaboration, Phys. Rev. Lett.**97**, 021802 (2006).
- [3] A. Abulencia *et al.*, CDF Collaboration, Phys. Rev. Lett. **97**,242003, (2006).
- [4] V.M. Abazov *et al.*, D0 Collaboration, Nucl. Instrum. and Meth. A**565** 463-537 (2006).
- [5] D0 Collaboration, Conference Note 5474.
- [6] V.M. Abazov *et al.*, Phys. Rev. D**74**, 112002 (2006).
- [7] D0 Conference Note 5210.
- [8] H.G. Moser, A. Roussarie, Nucl. Instrum. Meth. A**384** 491 (1997)